

Loss of butyrate-induced apoptosis in human hepatoma cell lines HCC-M and HCC-T having substantial Bcl-2 expression. *Hepatology*. 1998;27(5):1233-40. Hidetsugu Saito, Hirotooshi Ebinuma, Masahiko Takahashi, Fumihiko Kaneko, Kanji Wakabayashi, Mitsuyasu Nakamura, Hiromasa Ishii.

Antisense oligodeoxynucleotide against c-myc mRNA induces differentiation of human hepatocellular carcinoma cells. *Int. J. Oncology*. 1999;15:991-999. Ebinuma Hirotooshi, Saito Hidetsugu, Saito Yoshimasa, Wakabayashi Kanji, Nakamura Mitsuyasu, Kurose Iwao, Ishii Hiromasa

Bcl-2 related proteins are dramatically induced at the early stage of differentiation in human liver cancer cells by a histone deacetylase inhibitor projecting an anti-apoptotic role during this period. *Oncology Reports*.2000;7(2):285-288. Wakabayashi Kanji, Saito Hidetsugu, Ebinuma Hirotooshi, Saito Yoshimasa, Takagi Tamako, Nakamura Mitsuyasu, Umezawa Akihiro, Hata Junichi, Ishii Hiromasa

Reduction of telomerase activity in human liver cancer cells by a histone deacetylase inhibitor. *J Cell Physiol*. 2001;187(3):392-401. Mitsuyasu Nakamura, Hidetsugu Saito, Hirotooshi Ebinuma, Kanji Wakabayashi, Yoshimasa Saito, Tamako Takagi, Nobuhiro Nakamoto, Hiromasa Ishii

Interferon-associated retinopathy in a uniform regimen of natural interferon- α therapy for chronic hepatitis C Liver.2001;21(3):192-197 Saito Hidetsugu, Ebinuma Hirotooshi, Nagata Hiroshi, Inagaki Yasutaka, Nakamoto Nobuhiro, Saito Yoshimasa, Wakabayashi Kanji, Takagi Tamako, Nakamura Mitsuyasu, Katsura H, Oguchi Y, Ishii Hiromasa

Effect of the molar ratio of branched-chain to aromatic amino acids on growth and albumin mRNA expression of human liver cancer cell lines in a serum-free medium. *Nutr Cancer*. 2001;39(1):126-131. Saito Yoshimasa, Saito Hidetsugu, Nakamura Mitsuyasu, Wakabayashi Kanji, Takagi Tamako, Ebinuma Hirotooshi, Ishii Hiromasa

Interferon therapy for hepatocellular carcinoma patients with low HCV-RNA levels. *Hepato-gastroenterology*.2002;49(45):724-9. Miyaguchi S, Watanabe T, Takahashi H, Nakamura Mitsuyasu, Saito Hidetsugu, Ishii Hiromasa

The detection of IRF-1 promoter polymorphisms and their possible contribution to T helper response in chronic hepatitis C. *J Interferon Cytokine Res.* 2002;22(6):693-700
Saito Hidetsugu, Tada Shinnichirou, Wakabayashi Kanji, Nakamoto Nobuhiro, Takahashi Masahiko, Nakamura Mitsuyasu, Ebinuma Hirotohi, Ishii Hiromasa

Immune evasion by hepatitis C virus NS3/4A protease-mediated cleavage of the Toll-like receptor 3 adaptor protein TRIF. *Proc Natl Acad Sci USA.*2005;102(8):2992-7.
Kui Li, Eileen Foy, Josephine C. Ferreon, Mitsuyasu Nakamura, Allan C. M. Ferreon, Masanori Ikeda, Stuart C. Ray, Michael Gale, Jr., and Stanley M. Lemon

Association of pseudomembranous colitis with Henoch-Schonlein purpura. *J Gastroenterol.*2005;40:641-5
Takehiko Hayakawa, Hiroyuki Imaeda, Mitsuyasu Nakamura, Shunsuke Komoto, Kazuo Maruta, Hidetoshi Shiozu, Haruhiko Ogata, Yasushi Iwao, Hiromasa Ishii, Toshifumi Hibi

Down-regulation of the retinoblastoma tumor suppressor by the hepatitis C virus NS5B RNA-dependent RNA polymerase. *Proc Natl Acad Sci U S A.*2005;102(50):18159-64
Tsubasa Munakata, Mitsuyasu Nakamura, Yuqiong Liang, Kui Li, and Stanley M. Lemon

Possible Molecular Mechanism of the Relationship Between NS5B Polymorphisms and Early Clearance of Hepatitis C Virus During Interferon Plus Ribavirin Treatment
J.Med.Virol. 2008;80 (4):632-9.Mitsuyasu Nakamura, Hidetsugu Saito, Masanori Ikeda, Shinichiro Tada, Naoki Kumagai Nobuyuki Kato, Kunitada Shimotohno, Toshifumi Hibi

Advances in genomic research on hepatitis C virus with a useful tool, replicon system
*Keio J Med.*2008;57(2):75-83. Mitsuyasu Nakamura, Hidetsugu Saito, Toshifumi Hibi

Omega-3 fatty acids exacerbate DSS-induced colitis through decreased adiponectin in colonic subepithelial myofibroblasts. *Inflamm Bowel Dis* 2008;14(10):1348-57.
Matsunaga H, Hokari R, Kurihara C, Okada Y, Takebayashi K, Okudaira K, Watanabe C, Komoto S, Nakamura M, Tsuzuki Y, Kawaguchi A, Nagao S, Itoh K, Miura S.

Omega-3 polyunsaturated fatty acids ameliorate the severity of ileitis in the senescence accelerated mice (SAM)P1/Yit mice model. *Clini Exp Immunol.*2009;158(3)325-33

Matsunaga H, Hokari R, Kurihara C, Okada Y, Takebayashi K, Okudaira K, Watanabe C, Komoto S, Nakamura M, Tsuzuki Y, Kawaguchi A, Nagao S, Miura S.

Cilostazol, a specific PDE-3 inhibitor, ameliorates chronic ileitis via suppression of interaction of platelets with monocytes. *Am J Physiol Gastrointest Liver Physiol*.2009;297(6):1077-84. Matsunaga H, Hokari R, Higashiyama M, Kurihara C, Okada Y, Watanabe C, Komoto S, Nakamura M, Kawaguchi A, Nagao S, Miura S.

An antioxidant resveratrol significantly enhanced replication of hepatitis C virus. *World J Gastroenterol*. 2010;16(2):184-92. Mitsuyasu Nakamura, Hidetsugu Saito, Masanori Ikeda, Ryota Hokari, Nobuyuki Kato, Toshifumi Hibi, Soichiro Miura

Interferon-alpha increases monocyte migration via platelet-monocyte interaction in murine intestinal microvessel *Clini Exp immunol*. 2010;162(1):156-62. Higashiyama M, Hokari R, Kurihara C, Ueda T, Nakamura M, Komoto S, Okada Y, Watanabe C, Kawaguchi A, Nagao S, Miura S

Increased expression of lipocain-type-prostaglandin Dsynthase in ulcerative colitis and exacerbating role in murine colitis. *Am J Physiol Gastrointest Liver Physiol* 2011;300(3): G401-8. Hokari R, Kurihara C, Nagata N, AriakeK, OkadaY, WatanabeC, KomotoS, Nakamura M, KawaguchiA, NagaoS, Urade Y Miura S

Physiological stress exacerbates murine colitis by enhancing proinflammatorycytokine expression that is dependent on IL-18. *Am J Physiol Gsatrointest Liver Physiol* 2011; 301(3): G555-64.Matsunaga H, Hokari R, UedaT, Kurihara C, HozumiH, HigashiyamaM, OkadaY, WatanabeC, KomotoS, Nakamura M, Kawaguchi A., NagaoS, SekiyamaA, MiuraS,

Effects of Astaxanthin on Intestinal Microflora in Mice Fed a High-fat Diet. *Anti-Aging Medicine*.2013;10(4):77-91.Yoshikazu Yonei, Masayuki Yagi, Mitsuyasu Nakamura, Lanny Parengkuan, Mari Ogura, Toshio Taira, Sayori Asano , Hui-Hsiung Liu.